

# Reducing Nutrient Loss: Science Shows What Works



Iowa has been working for decades to protect and improve water quality. However, progress measured toward reduction targets at the watershed scale has been challenging, and many complex nutrient-related impacts in Iowa's lakes, reservoirs, and streams remain to be addressed.

The Iowa Nutrient Reduction Strategy is a science and technology-based framework to assess and reduce nutrients to Iowa waters and the Gulf of Mexico. It directs efforts to reduce nutrients in surface water from both point and nonpoint sources in a scientific, reasonable, and cost-effective manner.

It was prompted by the 2008 Gulf Hypoxia Action Plan that calls for Iowa and other states along the Mississippi River to develop strategies to reduce nutrient loadings to the Gulf of Mexico. The plan established a goal of at least a 45 percent reduction in total nitrogen and total phosphorus loads.

The Iowa strategy is a coordinated approach for reducing nutrient loads discharged from the state's largest wastewater treatment plants, in combination with targeted practices designed to reduce loads from nonpoint sources such as agriculture.

Success can be achieved using the tools known to work, such as targeted, voluntary conservation measures, in conjunction with research, development, and demonstration of new approaches. The goal is application of proven practices in fields and cities across Iowa.

## Science Provides Guidance

The strategy related to farmland is built on a scientific assessment of practices and associated costs to reduce loading of nitrogen (N) and phosphorus (P) to Iowa surface waters.

The College of Agriculture and Life Sciences at Iowa State University and the Iowa Department of Agriculture and Land Stewardship partnered to conduct the scientific assessment. The science team consisted of 23 individuals representing five agencies or organizations.

The objective of the science assessment was to identify and model the effectiveness of specific practices at reducing N and P reaching the Gulf of Mexico.

The assessment involved establishing baseline conditions, reviewing scientific literature, estimating potential load

reductions, and estimating implementation costs. The assessment shows that broad implementation of a combination of practices will be needed to reach desired load reductions.

## A Closer Look

The need to increase voluntary efforts to reduce nutrient loss is one of the key points related to agriculture in Iowa's Nutrient Reduction Strategy.

The science assessment identified effective nutrient reduction practices in three categories — nitrogen and phosphorus management, land use, and edge-of-field. (See charts on pages 3-4.)

**Management practices** involve application rate, timing, and method, plus the use of cover crops and reduced tillage.

**Land use practices** include perennial energy crops, extended rotations, grazed pastures, and land retirement.

**Edge-of-field practices** involve drainage water management, wetlands, bioreactors, buffers, terraces, and sediment control.

Some practices that have the greatest potential are highlighted here.

## Management Practices – Nitrogen

**Rate Reduction:** Matching N application rates with the Corn Nitrogen Rate Calculator, a university developed online tool, has potential to reduce nitrate-N loss. This tool estimates optimal N rates based on fertilizer and corn prices. (Find the calculator here: <http://extension.agron.iastate.edu/soilfertility/nrate.aspx>).

**Nitrification Inhibitor:** Research shows a corn yield increase plus a nitrate-N loss decrease when using a nitrification inhibitor (Nitrapyrin) with fall applied anhydrous ammonia. The only cost associated with this practice is the material. There is a corn yield increase of approximately 6 percent.

**Sidedress:** Sidedressing N can be done in different ways and with different sources of N, yet the concept of applying fertilizer after corn emergence is consistent. This strategy includes applying N during plant uptake, as well as timing to reduce the risk of loss from leaching events. Sidedressing also allows the N rate to be optimized by either soil sampling or crop canopy sensing.



## Management Practices – Phosphorus

**Consider Soil-Test P:** This practice involves not applying P on fields where the Soil-Test P (STP) values exceed the upper boundary of the optimum level for corn and soybean in Iowa. The practice would continue until the STP level reaches the optimum level.


**Cover Crops:** Planting a late summer or early fall seeded cover crop can reduce P loss. For example, winter rye offers benefits of easy establishment, seeding aerially or by drilling, growth in cool conditions, initial growth when planted in the fall, and continued growth in the spring. Cover crops also are effective at reducing N loss.



**Reduced Tillage:** Conservation tillage, where 30% or more of the soil surface is covered with crop residue after planting, or no-till, where 70% or more of the soil surface is covered with crop residue after planting, reduces soil erosion and surface runoff. Reduced erosion and runoff also reduces P transport.

## Land Use Practices – Nitrogen and Phosphorus

**Extended Rotations:** Extended rotations reduce the application and the loss of both P and nitrate-N. If a shift to extended rotations is significant, the amount of corn and soybean produced in Iowa would be reduced, along with an increase in alfalfa production that could support increased livestock production for alfalfa feeding. Another benefit would be improved soil quality.



**Energy Crops Replacing Row Crops:** Although there is not a current large market for perennial biomass crops as a source for energy or transportation fuel production, there are local and regional markets. Replacing row crops with energy crops or integrating energy crops within the rowcrop landscape decreases erosion, surface runoff, and leaching losses in the area implemented; therefore, the loss of both P and nitrate-N is reduced. An added benefit is an increase in wildlife habitat.

## Edge-of-Field Practices – Nitrogen and Phosphorus

**Wetlands:** Wetlands targeted for water quality benefits show great potential for nitrate-N reduction. Wetland costs include design, construction, buffer seeding, maintenance, and land acquisition. In addition to water quality benefits, these wetlands provide other benefits such as improved aesthetics and habitat.



**Bioreactors:** Subsurface drainage bioreactors also show good potential for nitrate-N reduction. Bioreactor costs include control structures, woodchips, design, construction, seeding, additional tile, management, and maintenance.

**Buffers:** Edge-of-field technologies such as buffers are designed to settle sediment and sediment-bound N and P, along with retaining nitrate-N and dissolved P. Buffers also provide wildlife habitat, sequester carbon, reduce greenhouse gas emissions, stabilize stream banks, and potentially reduce flood impacts. Costs of buffers can vary greatly depending on width, type of vegetation, and the amount of earthwork required.

## What's Next?

Iowa's Nutrient Reduction Strategy is a key step toward improving Iowa's water quality while ensuring the state's continued economic growth and prosperity. From it, operational plans are being developed through the Water Resources Coordinating Council. The strategy document will evolve over time as new information, data, and science are discovered and adopted.

The path forward to reducing nutrient impacts will not be easy, as it will require a high adoption rate of multiple practices to achieve the goal of cleaner water and a profitable agriculture. To learn more about the practices that may be right for your farm, attend a field day, contact the Iowa Department of Agriculture and Land Stewardship, or Iowa State University Extension and Outreach, or a certified crop adviser.

More information on Iowa's Nutrient Reduction Strategy is available at <http://www.nutrientstrategy.iastate.edu>.

# Iowa Strategy to Reduce Nutrient Loss: Nitrogen Practices

This table lists practices with the largest potential impact on nitrate-N concentration reduction (except where noted). Corn yield impacts associated with each practice also are shown as some practices may be detrimental to corn production. If using a combination of practices, the reductions are not additive. Reductions are field level results that may be expected where practice is applicable and implemented.

	Practice	Comments	% Nitrate-N Reduction <sup>+</sup>	% Corn Yield Change <sup>++</sup>
			Average (SD*)	Average (SD*)
Nitrogen Management	Timing	Moving from fall to spring pre-plant application	6 (25)	4 (16)
		Spring pre-plant/sidedress 40-60 split Compared to fall-applied	5 (28)	10 (7)
		Sidedress – Compared to pre-plant application	7 (37)	0 (3)
		Sidedress – Soil test based compared to pre-plant	4 (20)	13 (22)**
	Source	Liquid swine manure compared to spring-applied fertilizer	4 (11)	0 (13)
		Poultry manure compared to spring-applied fertilizer	-3 (20)	-2 (14)
	Nitrogen Application Rate	Nitrogen rate at the MRTN (0.10 N:corn price ratio) compared to current estimated application rate. (ISU Corn Nitrogen Rate Calculator – <a href="http://extension.agron.iastate.edu/soilfertility/nrate.aspx">http://extension.agron.iastate.edu/soilfertility/nrate.aspx</a> can be used to estimate MRTN but this would change Nitrate-N concentration reduction)	10	-1
	Nitrification Inhibitor	Nitrapyrin in fall – Compared to fall-applied without Nitrapyrin	9 (19)	6 (22)
	Cover Crops	Rye	31 (29)	-6 (7)
		Oat	28 (2)	-5 (1)
Living Mulches	e.g. Kura clover – Nitrate-N reduction from one site	41 (16)	-9 (32)	
Land Use	Perennial	Energy Crops – Compared to spring-applied fertilizer	72 (23)	
		Land Retirement (CRP) – Compared to spring-applied fertilizer	85 (9)	
	Extended Rotations	At least 2 years of alfalfa in a 4 or 5 year rotation	42 (12)	7 (7)
	Grazed Pastures	No pertinent information from Iowa – assume similar to CRP	85	
Edge-of-Field	Drainage Water Mgmt.	No impact on concentration	33 (32)	
	Shallow Drainage	No impact on concentration	32 (15)	
	Wetlands	Targeted water quality	52	
	Bioreactors		43 (21)	
	Buffers	Only for water that interacts with the active zone below the buffer. This would only be a fraction of all water that makes it to a stream.	91 (20)	

<sup>+</sup> A positive number is nitrate concentration or load reduction and a negative number is an increase.

<sup>++</sup> A positive corn yield change is increased yield and a negative number is decreased yield. Practices are not expected to affect soybean yield.

\* SD = standard deviation. Large SD relative to the average indicates highly variable results.

\*\* This increase in crop yield should be viewed with caution as the sidedress treatment from one of the main studies had 95 lb-N/acre for the pre-plant treatment but 110 lb-N/acre to 200 lb-N/acre for the sidedress with soil test treatment so the corn yield impact may be due to nitrogen application rate differences.



# Iowa Strategy to Reduce Nutrient Loss: Phosphorus Practices

Practices below have the largest potential impact on phosphorus load reduction. Corn yield impacts associated with each practice also are shown, since some practices may increase or decrease corn production. If using a combination of practices, the reductions are not additive. Reductions are field level results that may be expected where practice is applicable and implemented.

	Practice	Comments	% P Load Reduction <sup>a</sup>	% Corn Yield Change <sup>b</sup>
			Average (SD <sup>c</sup> )	Average (SD <sup>c</sup> )
Phosphorus Management Practices	Phosphorus Application	Applying P based on crop removal – Assuming optimal STP level and P incorporation	0.6 <sup>d</sup>	0
		Soil-Test P – No P applied until STP drops to optimum	17 <sup>e</sup>	0
	Source of Phosphorus	Liquid swine, dairy, and poultry manure compared to commercial fertilizer – Runoff shortly after application	46 (45)	-1 (13)
		Beef manure compared to commercial fertilizer – Runoff shortly after application	46 (96)	
	Placement of Phosphorus	Broadcast incorporated within 1 week compared to no incorporation, same tillage	36 (27)	0
		With seed or knifed bands compared to surface application, no incorporation	24 (46)	0
	Cover Crops	Winter rye	29 (37)	-6 (7)
	Tillage	Conservation till – chisel plowing compared to moldboard plowing	33 (49)	0 (6)
		No till compared to chisel plowing	90 (17)	-6 (8)
	Land Use Change	Perennial Vegetation	Energy Crops	34 (34)
Land Retirement (CRP)			75	
Grazed pastures			59 (42)	
Erosion Control and Edge-of-Field Practices	Terraces		77 (19)	
	Buffers		58 (32)	
	Control	Sedimentation basins or ponds	85	

<sup>a</sup> - A positive number is P load reduction and a negative number is increased P load.

<sup>b</sup> - A positive corn yield change is increased yield and a negative number is decreased yield. Practices are not expected to affect soybean yield.

<sup>c</sup> - SD = standard deviation. Large SD relative to the average indicates highly variable results.

<sup>d</sup> - Maximum and average estimated by comparing application of 200 and 125 kg P<sub>2</sub>O<sub>5</sub>/ha, respectively, to 58 kg P<sub>2</sub>O<sub>5</sub>/ha (corn-soybean rotation requirements) (Mallarino et al., 2002).

<sup>e</sup> - Maximum and average estimates based on reducing the average STP (Bray-1) of the two highest counties in Iowa and the statewide average STP (Mallarino et al., 2011a), respectively, to an optimum level of 20 ppm (Mallarino et al., 2002). Minimum value assumes soil is at the optimum level.

<sup>f</sup> - P retention in wetlands is highly variable and dependent upon such factors as hydrologic loading and P mass input.

## SPONSORS

**IOWA STATE UNIVERSITY**  
Extension and Outreach



Iowa State University Extension and Outreach programs are available to all without regard to race, color, age, religion, national origin, sexual orientation, gender identity, genetic information, sex, marital status, disability, or status as a U.S. veteran. Inquiries can be directed to the Director of Equal Opportunity and Compliance, 3280 Beardshear Hall, (515) 294-7612.